

贵州沟螺属一新种记述 (前鳃亚纲, 中腹足目, 环口螺科)

罗泰昌¹ 张卫红² 周卫川^{3*}

1. 贵州师范大学 生命科学院 贵阳 550001
2. 新疆大学生命科学与技术学院 乌鲁木齐 830046
3. 福建省出入境检验检疫局 福州 350001

摘 要 在整理贵州省荔波县茂兰喀斯特森林自然保护区陆生贝类标本时, 经鉴定发现 1 新种, 即茂兰沟螺 *Dioryx maolanensis* sp. nov., 文中对新种形态特征、栖息环境进行了描述, 并对其相似种进行了分析和讨论。
关键词 前鳃亚纲, 中腹足目, 环口螺科, 沟螺属, 新种。
中图分类号 Q959. 212

在整理贵州省荔波县茂兰喀斯特森林自然保护区陆生贝类标本时, 经鉴定发现环口螺科、沟螺属 1 新种, 即茂兰沟螺 *Dioryx maolanensis* sp. nov., 报道如下。
茂兰沟螺, 新种 *Dioryx maolanensis* sp. nov.
(图 1~ 6)

正模标本, 壳高 5. 16 mm, 壳宽 4. 16 mm, 壳口直径 1. 33 mm。副模标本 6 个, 壳高 4. 50~ 5. 33 mm, 壳宽 3. 5~ 5. 16 mm, 壳口直径 1. 00~ 1. 67 mm, 壳口宽 1. 33 mm, 2001 年 7 月 9 日采自贵州省荔波县茂兰石灰岩森林中。正、副模标本均保存在中国科学院动物研究所 (北京)。




图 1~ 2 茂兰沟螺, 新种 *Dioryx maolanensis* sp. nov.
1. 贝壳侧面观 (lateral view of shell) 2. 贝壳右侧面观 (right lateral view of shell)

贝壳小型, 右旋, 壳质厚, 坚实, 呈圆球形, 有 $3\frac{1}{2}$ 个螺层, 各螺层均膨胀, 体螺层特膨大, 但至壳口处紧缩, 呈 1 凹槽。壳顶钝, 胚螺层光滑, 缝合线深。壳面呈淡黄色, 陈旧标本呈灰白色, 并有稀疏而明显成纵行排列的近乎肋状条纹的生长线。

在体螺层背侧面靠近壳口处缝合线上有 1 细管状的呼吸管 (长 0. 83~ 1. 00 mm)。壳口呈圆形, 口缘外折并延长, 形成约 0. 8 mm 的口颈部, 口唇厚, 外卷形成双唇, 其边缘锋利, 并与体螺层分离。厣较厚为角质, 黄褐色, 与壳口同大, 半透明, 有 3 层,

贵州省高校发展专项资金资助自然科学类重点科研项目 (黔教科 2004112) 福建省自然科学基金项目 (2009J0104) 资助。
* 通讯作者。
收稿日期: 2008-08-20, 修订日期: 2009-07-20.




图 3~ 6 茂兰河螺, 新种 *Dioryx maolanensis* sp. nov.
3. 颚片 (jaw) 4. 齿舌 (radula) 5. 齿舌放大 (radula enlarged) 6. 厝 (operculum)

内层无螺旋纹, 光滑, 外层有一核心, 并有明显的螺旋纹。脐孔小而深, 略被小轴缘遮盖, 呈缝隙状。颚片薄片状, 透明, 其表面布满鳞片状结构。齿舌呈带状, 角质, 有中央齿 1 枚, 其尖端有 7 枚小齿; 侧齿 2 枚, 其尖端有 5 枚小齿; 缘齿 2 枚, 其尖端有 4 枚小齿。

栖息环境 常生活在阴暗潮湿多石灰岩的山区和丘陵地带, 一般栖息于石灰岩多腐殖质的缝隙、

松软潮湿的土壤表面, 落叶、腐木或石块下。
讨论 新种与乳阳沟螺 *Dioryx nuyangensis* Hu, Yin and Chen, 2004 相近似, 但新种贝壳较小 (壳高 5.16 mm, 壳宽 4.16 mm, 壳口直径 1.33 mm), 有 3 $\frac{1}{2}$ 个螺层, 呼吸管长 0.83~ 1.00 mm, 壳面有稀疏而明显成纵行排列的近乎肋状条纹和生长线, 颚片薄片状, 透明, 其表面布满鳞片状结构。齿舌呈带状,

角质, 有中央齿 1 枚, 其尖端有 7 枚小齿, 侧齿 2 枚, 其尖端有 5 枚小齿, 缘齿 2 枚, 其尖端有 4 枚小齿。后者个体较大 (壳高 7.10 mm, 壳宽 5.50 mm, 壳口直径 1.75 mm), 有 $4 \sim 4\frac{1}{2}$ 个螺层, 呼吸管长 2.80 mm。壳面无成纵行排列的近乎肋状条纹和生长线。故有所区别。

REFERENCES (参考文献)

- Benson, W. H. 1859. A sectional distribution of the genus *Alycaeus* Gray, with characters of 6 new species and of other Cyclostoniidae, collected at Darjiling by W. T. Blanford, Esq., *Ann. Mag. Nat. Hist. Lond.*, 3 (3): 176-184.
Chen, D N and Gao, J X 1987. Economic Fauna Sinica of China

- (Terrestrial, Mollusca). Science Press, Beijing. 187. [陈德牛, 高家祥, 1987. 中国经济动物志, 陆生软体动物. 北京: 科学出版社. 1~187]
Chen, D N and Zhang, G Q 2001. Five new species of land snails from the Tibet Autonomous Region, China. *Acta Zootaxonomica Sinica*, 26 (1): 11-17. [陈德牛, 张国庆, 2001. 西藏陆生贝类五新种记述. 动物分类学报, 26 (1): 11~17]
Hu, Z Q, Yin, H Q and Chen, D N 2004. A new species of the genus *Dioryx* Benson from China (Prosobranchia, Mesogastropoda, Cyclophoridae). *Acta Zootaxonomica Sinica*, 29 (4): 704-705. [胡自强, 银海强, 陈德牛, 2004. 中国沟螺属一新种记述 (前鳃亚纲, 中腹足目, 环口螺科). 动物分类学报, 29 (4): 704-705]
Heude, P. M. 1882-1885. Notes sur les mollusques terrestres de la vallée du Fleuve Bleu, *Mem. D'Hist. Nat. l'Emp. Chinois.*, 97-98. pl. 24, figs. 6, 6a, 6b, 6c-7.
Yen, T. C. 1939. Die Chinesisch Land und Süßwasser Gastropoden des Natur Museum Senckenberg. *Abh. Senckenb. Natur. Ges.*, 444: 28-29, pl. 2, f. 28-30.

A NEW SPECIES OF THE GENUS *DIORYX* BENSON FROM CHINA (PROSOBRANCHIA, MESOGASTROPODA, CYCLOPHORIDAE)

LUO Tai Chang¹, ZHANG Wei Hong², ZHOU Wei Chuan^{3*}

1. School of Life Sciences, Guizhou Normal University, Guiyang 550001, China
2. College of Life Science and Technology, Xinjiang University, Urumqi 830046, China
3. Fujian Entry-Exit Inspection & Quarantine Bureau, Fuzhou 350001, China

Abstract This paper deals with a new species of the genus *Dioryx* collected from Guizhou Province, China. Type specimens are deposited in the Institute of Zoology, Chinese Academy of Science, China.

Dioryx maolanensis sp. nov. (Figs. 1-6)

Holotype, alt. 5.16 mm, diam. 4.16 mm, diam. of aperture 1.33 mm. Paratypes 6 specimens, alt. 4.50-5.33 mm, diam. 3.5-5.16 mm, diam. of aperture 1.00-1.67 mm, collected from Madan Town, Libo County (25°03' N, 108°00' E), Guizhou Province, China, 9 July 2001.

Shell small, dextral, thick, globose conoid. Whorls $3\frac{1}{2}$, moderately convex, spire high, body whorl very convex. Shell yellowish in colour, worn specimens ashen white, surface with very distinct vertically thin ribbed line and growth line, protoconch smooth, without spiral line. Apex blunt. Suture deep, with a 0.83-1.00 mm wormform projection at suture of body whorl. Aperture circular, thick, with two lips, somewhat expanded and reflexed. Operculum circular, cuticular, yellow brown in colour, and the inner surface has papuleform projection, and concentric circles lines indistinct visible in the surface. Umbilicus small, crevice

formed. Jaw thin flake-shaped, transparent, the surface with scale shaped structure. The radula tape shaped, circular, with 1 central tooth, and with 7 smaller cusps on apex; 2 lateral teeth, and with 5 smaller cusps on apex; 2 marginal teeth, and with 4 smaller cusps on apex.

This new species is similar to *Dioryx nuyangensis* Hu, Yin and Chen, 2004, but differs obviously from the latter in the small size; the shell with whorls $3\frac{1}{2}$, length 5.16 mm, diam. 4.16 mm; diam of aperture 1.33 mm; surface of shell with very distinct vertically thin ribbed line and growth line, with a 0.83-1.00 mm wormform projection at suture of body whorl. Jaw thin flake shaped, transparent, the surface with scale shaped structure. The radula tape shaped, circular, with 1 central tooth, and with 7 smaller cusps on apex; 2 lateral teeth, and with 5 smaller cusps on apex; 2 marginal teeth, and with 4 smaller cusps on apex. The shell of the latter species with whorls $4\frac{1}{2}$, length 7.10 mm, diam. 5.50 mm; diam of aperture 1.75 mm; with a 2.80 mm wormform projection at suture of body whorl.

Key words Prosobranchia, Mesogastropoda, Cyclophoridae, *Dioryx*, new species.

* Corresponding author.